

Year in Review: Clinical Investigator Perspectives on the Most Relevant New Datasets and Advances in Bruton Tyrosine Kinase Inhibitors for Chronic Lymphocytic Leukemia

THE CORRECT ANSWER IS INDICATED WITH YELLOW HIGHLIGHTING.

- 1. Which of the following descriptions best characterizes the study design of the BRUIN CLL-314 trial?**
  - a. A Phase II study evaluating time-limited pirtobrutinib for patients with chronic lymphocytic leukemia (CLL) that was previously treated with prior Bruton tyrosine kinase (BTK) inhibitor and Bcl-2 inhibitor therapy
  - b. A Phase III superiority study evaluating pirtobrutinib versus ibrutinib for relapsed/refractory CLL
  - c. A Phase III noninferiority study evaluating pirtobrutinib versus ibrutinib for patients with CLL that was treatment-naïve or previously treated with non-BTK inhibitor therapy**
- 2. Which of the following outcomes best describes the reduction in risk of disease progression or death in the Phase III BRUIN CLL-321 study evaluating pirtobrutinib monotherapy versus idelalisib/rituximab or bendamustine/rituximab for relapsed/refractory CLL?**
  - a. Pirtobrutinib led to a significant 28% reduction
  - b. Pirtobrutinib led to a significant 46% reduction**
  - c. Pirtobrutinib did not lead to a significant reduction in the risk of disease progression or death
- 3. Recently the FDA granted approval to acalabrutinib for which of the following indications in adult patients with CLL/ small lymphocytic lymphoma (SLL)?**
  - a. As monotherapy in the first-line setting
  - b. As monotherapy in the relapsed/refractory setting for patients who received prior treatment using a covalent BTK inhibitor
  - c. As combination therapy with venetoclax in the first-line setting**
  - d. As combination therapy with venetoclax in the relapsed/refractory setting for patients who received prior treatment using a covalent BTK inhibitor
- 4. Which of the following Grade  $\geq 3$  adverse events was most commonly reported among patients receiving zanubrutinib, obinutuzumab and sonotoclax for previously treated CLL in the Phase I BGB-11417-101 study?**
  - a. Febrile neutropenia
  - b. Neutropenia**
  - c. Infusion-related reaction
- 5. Arm D of the Phase III SEQUOIA study demonstrated promising efficacy with the combination of zanubrutinib and venetoclax in which population of patients with treatment-naïve CLL?**
  - a. All-comers
  - b. Those with IGHV-unmutated disease
  - c. Those with del(17p) and/or TP53 mutations**
  - d. Those with BTK C481S mutations